



MiteGard®



SPECIMEN LABEL

CAN BE USED FOR ORGANIC PRODUCTION

• Poultry & Livestock Lice & Mite control

This product meets the Program Standards set forth by the National Organic Program (NOP) final rule 7CFR Part 205.

Active Ingredients	
Rosemary	6.0%
Sesame	5.8%
Peppermint	5.4%
Soybean Oil	5.1%
Thyme	4.4%
Cinnamon	3.9%
Malic Acid	3.4%
Garlic	2.7%
Other Ingredients	
Beeswax, Calcium Carbonate, Carrageenan, Cellulose, Citric Acid, Glycerin, Kaolin, Lecithin, Ground Mustard, Palm Oil, Sodium Bicarbonate, Sodium Chloride	63.3%

This product has not been registered by the US Environmental Protection Agency. USAgritech, Inc. represents that this product qualifies for exemption from registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt, long pants, socks and shoes when mixing or using this product. Remove and wash contaminated clothing before reuse.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. If swallowed: Call a poison control center or doctor immediately for treatment advice. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

FOR USE AS A LIVESTOCK AND POULTRY SPRAY

MiteGard® is prepared using USAgritech's proprietary NEO Biological Blending Technology and is to be used as a contact livestock and poultry spray miticide and preventative pest agent for the control of mites on all animals and fowl.

DIRECTIONS FOR USE

General Information and Tips for Best Performance

For best results, buffer water pH to a neutral or acidic range (7.0pH or lower). Use outside of this recommended range may result in reduced performance on target pests. Application should be made as soon as mites appear. Apply directly to animals and fowl as well as surrounding areas for best control.

Method of Application for MiteGard®

Application using hand sprayers, mechanical sprayers/misters or dipping baths is recommended. Refer to Chart for specific application and dilution rates. For maximum control, apply MiteGard® directly on the animal's skin, hair or feathers and on all ground surfaces that contain insect eggs and larvae.

MiteGard® sprayed at regular intervals will help interrupt the insect development cycle thereby reducing insect population.

Dilution Instructions

MiteGard® is provided as a concentrate. Bag contents must be blended by shaking the bag prior to opening. After blending, fill a container with 25 gallons of water. Open bag by cutting off a top corner and SLOWLY empty contents into water while agitating water gently to blend. Stir solution lightly to fully blend. When blended, no further agitation is required. For best product performance use a pH range of 5.5 to 6.5. Use a plain citric acid buffer only.

Livestock & Poultry Dilution and Use Rates	Ounces of Concentrate per One Gallon of Water
For most hand and mechanical sprayer applicators	
For control of mites on cattle, horses, goats and hogs	0.75 fl. oz.
For Dipping Baths	0.5 fl. oz. to 0.75 fl. oz.
For effective control of Biting and Sucking Lice on cattle, goats and hogs	0.75 fl. oz.
To control poultry lice	0.5 fl. oz.
For control of mites on poultry	0.5 fl. oz.

Livestock and Poultry Mites including but not limited to: Chicken Mites (Red Roost Mite), Chiggers, Dust Mite, Lice (Biting and Sucking), Poultry Lice, Mange Mite, Northern Fowl Mite, Red Chicken Mite. MiteGard® is a general use contact miticide that is shown to control mite populations of most species of mites. This product has proven to kill adults, immatures and eggs. Apply as needed.

Application Timing for MiteGard®

Begin spraying at first sign of pest. Repeat spray on a 3 to 4 week interval depending on pest pressure.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present. Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

Storage

Store in original UV protected packages only. Store containers in a cool, dry location, away from direct sunlight, sources of intense heat or where freezing is possible. In case of spill on floor or paved surfaces, hose off with clean water. Store at temperatures between 41°F to 85°F (5°C to 30°C).

Disposal

Wastes resulting from using this product may be disposed of on site. Dispose of empty bag in a sanitary landfill or as permitted by state and local authorities

CONDITIONS OF SALE AND WARRANTY

USAgritech, Inc. warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. USAgritech, Inc. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR USE. Critical and unforeseeable factors beyond USAgritech's, Inc. control prevent it from eliminating all risks in connection with the use of this product. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather conditions, presence of other materials, drift to other crops or property, the manner of use or application, or failure to follow label directions will be assumed by the Buyer or User. Handling, storage and use of the product by Buyer and User are beyond the control of USAgritech, Inc. In no case shall USAgritech, Inc. be held liable for consequential, special or indirect damages resulting from the use of this product. The limit of USAgritech, Inc. liability shall be the purchase price for the quantity involved.

MiteGard is a registered Trademark of USAgritech, Inc.
Copyright USAgritech, Inc., 2009

Net Contents - 26 Fl Oz (769ml)

041409

